



REC'D 1 6 DEC 2003

The Patent Office Concept House Cardiff Road Newport South Wales NP10 8QQ

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely substantial company to certain additional company law rules.

Signed

Dated

25 November 2003

PRIORITY DOCUMENT


lahoney

SUBMITTED OR TRANSMITTED IN COMPLIANCE WITH RULE 17.1(a) OR (b)

BEST AVAILABLE COPY

Patents Act 1977 (Rule 16)





<u> 04NDVO2 E760447-2 D02094</u>

Request for grant of a patent



The Patent Office

Cardiff Road Newport Gwent NP9 1RH

Your reference 1.

P01/7700 0.00-0225503.2 2. Patent application number 0225503.2 **Zoolife International Limited** Full name, address and post code of the or 3. 55 Elizabeth Street each applicant London SW1W 9PP 8445181001

Patents ADP number

If the applicant is a corporate body, give the country/state of its incorporation

United Kingdom

4. Title of the invention **GEL FEED**

Name of your agent 5.

VENNER, SHIPLEY & CO

"Address for service" in the United Kingdom to which all correspondence should be sent

20 LITTLE BRITAIN LONDON EC1A 7DH

Patents ADP

1669004

6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or each of these earlier applications and the or each application number Country

Priority application number

Date of filing

7. If this application is divided or otherwise derived from an earlier UK application, give the number and filing date of the earlier application

Number of earlier application

Date of Filing

8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer 'YES' if:

- a) any applicant in 3. above is not an inventor, or b) there is an inventor who is not named as an applicant, or
- c) any named applicant is a corporate body)

YES

Form 1/77

11.

Enter the number of sheets for any of the 9. following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form

Description 10

> 2 Claim(s)

Abstract

Drawing(s)

If you are also filing any of the following state 10. how many against each item.

Priority documents

Translations of priority documents

Statement of inventorship and right to grant of a patent (Patents Form 7/77)

> Request for preliminary examination and search (Patents Form 9/77)

> > Request for substantive examination (Patents Form 10/77)

> > > Any other documents

I/We request the grant of a patent on the basis of this application.

Signature

1 November 2002

Name and daytime telephone number of 12. person to contact in the United Kingdom SIAN HUGHES 020 7600 4212

Gel Feed

5

10

25

The present invention relates to improvements in animal feed compositions, and in particular, to improvements in feed for aquatic animals, especially those kept in aquaria.

Food harvesting, processing, manufacture and storage may lead to a reduction in the nutritional value of the feed ingredients. Exposure to light, heat, pressure, mechanical actions, atmospheric conditions or irradiation may damage feed ingredients resulting in reduced quantities of nutrients and/or reduced bioavailability of important dietary components. Affected nutrients may include fat and water soluble vitamins, fat and water soluble carotenoids, immunomodulators and bioflavenoids.

As a result of this, the diets of captive species may not adequately reflect the nutritional diversity and bioavailability of the natural diet for a given species. This may result in nutritional restrictions and subsequent negative effects on growth, reproduction, health and sustainability of captive populations. This is of particular significance to species conservation programmes and in many areas of zoo and wildlife nutrition.

Carotenoids are pigments of plant origin and are known to act as powerful antioxidants. Certain carotenoids are additionally known to provide pigmentation and coloration of animal tissues. They are considered to be an important dietary component for many species.

Diets may require supplementation with additional amino acids to overcome amino acid restriction and to aid the provision of balanced protein within the ration.

It is known to add dry vitamin, mineral, carotenoid and amino acid powdered preparations to enrich feed. However, this has a number of disadvantages.

Distribution of such powdered particles or fine aggregates on or within the feed is affected by factors such as feed particle size and adherence. Post enrichment

-2-

settlement may occur, providing variance in feed quality, especially following storage, transport and distribution.

Liquid preparations containing vitamins, minerals, carotenoids and/or amino acids simplify enrichment and provide enhanced uniform distribution and adherence to the feed. In many cases the liquid may penetrate the feed surface and become strongly bound to the feed matrix.

In some formulations of the diet, a separate liquid supplement including a number of components in a liquid state may be supplied for addition to the feed at the time of mixing. Such liquids may include, but are not restricted to, amino acids, glucans, vitamins, minerals, carotenoids and oils/fats, all in a liquid form as an emulsion, solution or suspension.

The present invention seeks to address the problems or disadvantages associated with the known feed additives. The compositions according to the invention are preferably formulated to allow the simultaneous addition to feeds of highly bioavailable components in a liquid state, namely one or more of the following:

- 20 1. water soluble vitamins;
 - 2. water soluble minerals;
 - 3. aqueous soluble 1,3 /1,6 beta glucans;
 - 4. fat soluble vitamins in micelle form;
 - 5. water soluble carotenoids;
 - 6. fat soluble carotenoids in micelle form;
 - 7. amino acids in the liquid phase which may be presented in the oil or water phase of the composition, the presentation of the amino acids in the oil phase will be in micelle form; and
 - 8. a balanced mixture of oils.

30

25

5

The provision of water soluble vitamins in solution, fat soluble vitamins in micelles and amino acids in a liquid state in micelle and/or the aqueous phase provides a high degree of bioavailability and reduces the potential for chemical interactions

which may reduce bioavailability following consumption. A balanced addition of oils of specific chain length may be used to further aid enrichment.

It has been shown, for example, that the use of a liquid carotenoid supplement presenting the carotenoid in micelle form in an emulsion has a high bioavailability and is able to result in change in skin coloration in animals, particularly in fish and reptiles. The performance of such emulsions may exceed that of dry formulations.

A further advantage is that the composition of the liquid additive product continues to protect the vitamins and carotenoids against oxidation in the post application stage.

Animals may be fed gel diets varying in formulation and hardness. To date, gel diet formulations have had a number of inherent disadvantages.

The known gel diet formulations are supplied as a dry powder – this has potential problems with regard to settlement and distribution as set out above.

The known gel diet formulations also contain single or multiple binders as feed additives. Such use may reduce the value of the diet by: (1) reducing the nutrient density of the food; (2) affecting diet digestibility and adsorption/absorption following consumption, for example, some carbohydrates are known to reduce protein uptake and utilization in fish, particularly in cold water marine fish; and (3) changing the nutrient profile of the diet.

Yet another disadvantage associated with known gel diets is that there is a frequent requirement to use heat in order to activate the gelling process. Hot or boiling water or steam is often recommended for reconstitution of the powder into a gel. This may cause degradation of delicate nutrients, reducing nutrient availability and altering the nutritional value of the food. Digestibility may be altered, and this may have undesirable effects in terms of health, growth and condition of animals.

15

20

10

25

The amounts of gel binders used in conventional gel diets will vary, but is frequently high in cases where a high gel strength is required.

Where cold gel setting processes have been used for the production of diets, the process involves the addition to the diet of synergistic gelling agents to vary the speed of gel setting. Such additives may include organic and inorganic compounds.

According to a first aspect of the present invention, a composition is provided which comprises a dry powdered diet which, when mixed with wet feed and/or with water, forms a gel.

In a preferred embodiment of the invention, certain components of the diet may be supplied in a liquid state, as described above.

The water used for forming the gel may vary in formulation and salinity and is preferably cold.

In a preferred embodiment, the composition comprises amino acids, glucans, vitamins, minerals, peptides, nucleotides, bioflavinoids, oils, polysaccharides carotenoids and/or lipids.

In another preferred embodiment, the composition according to the first aspect of the invention contains ingredients which have natural binding properties, for example particulate natural algae.

When used as a natural binder in the composition of the present invention, the particulate natural algae particle size may vary, and is preferably micronised to a size of between 30 micron and 60 micron in diameter. Algae may be derived from marine or freshwater sources, and may be naturally sized or ground following harvesting and drying. The use of algae in the feed composition provides natural binding properties by action of compounds such as carragenans.

25

30

20

- 5 -

Conventional binders, for example cellulose ethers such as hydroxyethylcellulose and methylcellulose, can be included in an amount up to 0.001%.

Powdered yeasts and algae may be subjected to processing prior to inclusion in the composition to enhance bioavailability of components. For example, the pretreatment of algaes can lead to an increase in the bioavailability of carotenoid pigments.

5

15

20

25

In a further preferred embodiment, the composition may include feed attractants such as Betaine where an increase in palatability is desired.

In a yet further embodiment of the present invention, naturally or artificially enriched crustaceans and plankton may be added to the composition. This may include, but is not restricted to, *Daphnia pulicaria* enriched with *Aphanizomenon flosaquae*, and particularly crustaceans dried by refractance window drying systems or freeze drying.

The composition of the present invention may also include a conventional vitamin and mineral premix.

Compositions are also proposed which include organic minerals and minerals in a chellated form, such as selenium-methionine, where the mineral is bound to an amino acid (for example selanomethionine, which for example may be present in some yeasts). The use of organic minerals may aid mineral absorption, tissue retention and bioavailability.

Antioxidants and preservatives may be added to the composition to confer stability during storage, manufacture and during the post manufacture stage.

Where the composition according to the present invention has a separate liquid component, this liquid component has an aqueous phase containing water soluble vitamins, minerals, carotenoids and amino acids. Fats and fat soluble vitamins, carotenoids and amino acids are contained in a micelle "microencapsulated" form

distributed evenly throughout the liquid component. Plankton and algae may also be included in liquid form. The liquid component preferably contains water, emulsifiers, antioxidants, and preservatives.

The formation of the micelle or "microencapsulated" fat soluble ingredients is achieved by the use of conventional emulsifying agents, such as Polysorbate 80, and by high speed blending or emulsification. The micelle structures will preferably contain hydrophobic fat soluble vitamins and carotenoids within a hydrophilic protective envelope structure.

10

15

30

Stabilization of the fat soluble phase and optimisation of micelle distribution is achieved by the inclusion of conventional stabilizing agents, such as monopropylene glycol. The use of such stabilizing agents reduces the potential for product turbidity and affords excellent product clarity.

Preservation of the solution is achieved by the use of conventional preservatives, such as phosphoric acid and potassium sorbate. This prevents the growth of bacteria, fungi and yeasts.

Antioxidants may also be added to the composition to aid stability. Examples of such antioxidants include ascorbyl polyphosphate and butylated hydroxy-toluene.

Antioxidants prevent or minimize the loss of the active components of the composition. The incorporation of antioxidants thereby extends the shelf life of the composition and provides protection to the finished product in the post application phase.

In a further embodiment of the invention, cellulose, gum and sugar derivatives are added to the liquid composition to aid dispersion within or onto feeds in certain applications by virtue of their ability to increase solution viscosity and adherence. These products are not required for emulsification in this composition.

The fat and water soluble vitamins used in the composition may be a single or multiple. The quantity and ratio of vitamins included will be dependent on the

desired vitamin content of the feed produced using the composition of the present invention.

Similarly, the type and quantity of a number of optional additives included in the composition will vary, depending upon the desired make-up of the resultant feed product. Such variable additives include fat and water soluble carotenoids, aqueous soluble beta 1,2/1,6 glucans, specific chain length oils, and amino acids.

In one embodiment of the invention, suspensions of glucans and immunomodulating agents including algae are used.

20

In another embodiment, the amino acids are added in a free form to aid bioavailability.

15 It has been found that the composition according to the first aspect of the present invention affords stability to the active components during storage, application and post application period.

The composition provides a uniform dispersion of the various components within the composition and this aids optimal distribution of the components when the composition is applied to feeds.

The presence of the uniform micelles aids the uptake of fats and fat soluble vitamins carotenoids and amino acids from the diet at the level of the digestive tract. This is combined with the simultaneous presentation of water soluble vitamins, carotenoids and amino acids providing excellent bioavailability. The presence of amino acids in a liquid free form provides a high degree of bioavailability without the requirement for hydrolysis.

Thus, it is clear that the gel feed produced using the composition according to the first aspect of the present invention affords a number of important advantages over the known gel feed.

_ 8 -

According to a second aspect of the present invention, a gel feed product is provided, comprising the composition according to the first aspect of the present invention, mixed with wet feed and/or with water.

This gel feed product may be used for a variety of dietary applications. The ratio of composition to wet feed or water can be varied, depending upon the desired end product.

This gel feed has a number of advantages. Firstly, its formation does not require
heat. This means that the additives that may be included in the gel feed will not be
denatured or spoilt. This also makes it easier to include medicines in the gel feed.

Secondly, this cold setting instant gel feed may be prepared rapidly and may be fed as a moist crumb, or may be compressed to form a block and the block may be cut into pieces of a size and shape which suits the species being fed. Alternatively, the gel may be pressed into feeders such as corals for marine fish. The gel may also be pressed onto rope or alternative feeders for presentation. The feed can also be presented in such a way that only a certain species in the water can or will access it, thereby allowing targeted feeding.

15

20

Thirdly, the easily manipulated gel can be pressed or extruded into the desired shape, which it will then maintain for a relatively long period as the gel feed is relatively stable in water and only gradually disintegrates.

The use of the gel feed within corals, on rope or in/on an alternative holder such as a mesh or bobbin can target feed animals and will also increase the period of food availability, reducing competitive feeding behaviour and allowing natural grazing.

This is particularly important when feeding fish in captivity.

In order that the present invention may be better understood, a specific example will now be described, by way of illustration only.

This specific example of a gel feed according to the present invention is an omnivore feed intended for marine fish.

The composition comprises the following components or ingredients:

	Fucus 60 micron	481.00g
	Nori micronised	3868.00g
	Dulse micronised	1040.00g
	Spirulina	204.00g
10	Fishvits premix	210.00g
	Nori fine flake	808.52g
	Shrimp meal	1.214kg
	Aphanizomenon flos-aquae	133.77g
	Beta glucan premix	428.75g
15	Phaffia	330.28g
	Daphnia pulicaria RF dried	1030.91g
	Soya isolates	247.91g
	Selanomethionine as yeast	3.00g

5

30

The above components were blended so as to form a homogeneous powder. Then the powder was mixed with defrosted or fresh wet fish feeds in a ratio of 50:50 by weight.

The mixture was then blended in a paddle mixer for a short period, preferably

between 10 seconds and 2 minutes in most cases. This led to the formation of gel
in the form of semi-moist "crumbs".

Water was then added at 10-25mls for each 100g of dry diet used in the formulation whilst mixing again for a short period (10 seconds to 2 minutes in most cases). This formed a gel in the form of moist "crumbs".

The crumb formed in that way may be fed immediately and may be formed by hand or by machine into blocks or balls. These blocks or balls may be cut with a knife

into pieces of a size and shape to suit the animals being fed. Alternatively the block may be pressed onto a feeder/rope and hung or placed within the water as a food source.

In many cases the gel lasts for several hours as a gel. Advantageously, there is very little clouding of the water or gel diet breakdown whilst the gel feed is in the water.

Furthermore, the gel sets instantly without the need for it to be chilled. This means that the gel is very easy to manipulate and handle. This is particularly useful when the gel is to be applied to ropes, meshes and the like, which would be difficult, if not impossible, if the gel needed to chill in order to set. It also means that the gel will not quickly "melt" at room temperature or warmer.

Claims

- 1. A composition comprising a dry powdered diet which, when mixed with wet feed and/or with water, forms a gel.
- 2. A composition as claimed in claim 1, further comprising a liquid diet component.
- 3. A composition as claimed in either of claims 1 and 2, wherein the diet comprises one or more of vitamins, minerals, 1,3 /1,6 beta glucans, carotenoids, amino acids and oils.
 - 4. A composition as claimed in any one of the preceding claims, wherein the composition includes a natural binder.
 - 5. A composition as claimed in claim 4, wherein the natural binder is particulate natural algae.
- 6. A composition as claimed in claim 2, wherein the liquid component has an aqueous phase containing water soluble vitamins, minerals, carotenoids and/or amino acids.
 - 7. A composition as claimed in either of claim 2 or 6, wherein the liquid component has micelles containing fats, fat soluble vitamins, carotenoids and/amino acids.
 - 8. A composition as claimed in any one of claims 2, 6 and 7, further comprising plankton and/or algae in liquid form.
- 9. A gel feed comprising a composition as claimed in any one of the preceding claims mixed with wet food and/or water.
 - 10. A gel feed formed on to a rope.

5

15

- 11. A gel feed as claimed in claim 9, wherein the gel feed is a gel feed as claimed in claim 9.
- 5 12. A method of forming a cold setting gel feed comprising mixing a composition as claimed in any one of claims 1-8 with a wet food and/or water.

Abstract

Gel Feed

The present invention provides a composition comprising a dry powdered diet which, when mixed with wet feed and/or with water, forms a gel. The composition may optionally include a liquid component. A gel feed comprising the composition is also provided.

This Page is Inserted by IFW Indexing and Scanning Operations and is not part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

□ BLACK BORDERS
□ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
□ FADED TEXT OR DRAWING
□ BLURRED OR ILLEGIBLE TEXT OR DRAWING
□ SKEWED/SLANTED IMAGES
□ COLOR OR BLACK AND WHITE PHOTOGRAPHS
□ GRAY SCALE DOCUMENTS
□ LINES OR MARKS ON ORIGINAL DOCUMENT
□ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

IMAGES ARE BEST AVAILABLE COPY.

OTHER:

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.